

ORIGINAL ARTICLE

Historical Perspectives and Unani/Persian Concept of *Hijāma* (Cupping Therapy): A Time-Honored Remedy for Various Health Ailments

173

Abstract

Three different treatment modalities are detailed in the traditional textbook of Unani/Persian medicine, namely, '*Ilāj bi'l Tadbīr*' (Regimenal therapy), '*Ilāj bi'l Taghthiya*' (Dietotherapy), '*Ilāj bi'l Dawā*' (Pharmacotherapy), and '*Ilāj bi'l Yad*' (Surgery). This article explores *Hijāma* (cupping therapy) within the framework of the Unani/Persian system of medicine, highlighting its historical evolution, various types, therapeutic applications, mechanisms of action, and contraindications. The review aims to provide a thorough, comprehensive understanding of *Hijāma* and its significance in both traditional and contemporary healthcare. This review of *Hijāma* was conducted by searching classical Unani/Persian textbooks and major scientific databases, including PubMed, ScienceDirect, and Google Scholar, for additional information. *Hijāma* is an important regimen of *Ilāj bi'l Tadbīr* in the Unani system of medicine used for managing various diseases. It involves creating a partial vacuum in cups placed on the body using fire or suction to remove morbid materials, improve blood circulation, and divert harmful substances from affected tissues. The Unani emphasis on safe, minimally invasive therapies, combined with modern scientific approaches, offers promising prospects for future research and therapeutic applications.

Key words: *Hijāma*, Cupping Therapy, Unani Medicine, '*Ilāj bi'l Tadbīr*', Therapeutics, Pain Management, Delivery of Health Care

Received: 26 Jan 2025; Accepted: 11 Dec 2025; Online published: 31 May 2026
Research on History of Medicine/ 2026 May; 15(2): 173-184.

Khalid Rahim Wani (M.D.)¹ 
Abdul Nasir Ansari (M.D.)²
Mohd Nayab (M.D.)³

1- Assistant Professor, Department of '*Ilāj bi'l Tadbīr*' (Regimenal Therapy), Kashmir Tibbia College, Hospital and Research Centre, Taleemabad Shilvath, Bandipora, Jammu and Kashmir, India

2- Professor and Head, Department of '*Ilāj bi'l Tadbīr*' (Regimenal Therapy), National Institute of Unani Medicine, Kottigepalya, Magadi Main Road, Bengaluru-91, Karnataka, India

3- Associate Professor, Department of '*Ilāj bi'l Tadbīr*' (Regimenal Therapy), National Institute of Unani Medicine, Kottigepalya, Magadi Main Road, Bengaluru-91, Karnataka, India

Correspondence:

Khalid Rahim Wani
Assistant Professor, Department of '*Ilāj bi'l Tadbīr*' (Regimenal Therapy), Kashmir Tibbia College, Hospital and Research Centre, Taleemabad Shilvath, Bandipora, Jammu and Kashmir, India
e-mail: khalidwani4470@gmail.com

Citation:

Wani, Kh.R., Ansari, A.N., Nayab, M., 2026. Historical Perspectives and Unani/Persian Concept of *Hijāma* (Cupping Therapy): A Time-Honored Remedy for Various Health Ailments. *Res Hist Med*, 15(2), pp. 173-184. doi: 10.30476/rhm.2025.105684.1281.

Copyright: © Journal of Research on History of Medicine. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 Unported License, (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited non-commercially.

Introduction

Hijāma (cupping therapy) is considered one of the oldest therapeutic procedures in traditional medicine worldwide (Turk, and Allen, 2009). The precise origin of *Hijāma* remains controversial; however, historical evidence suggests it has been practiced for thousands of years across various ancient civilizations. Arabic historians report that the Assyrians were among the earliest communities to practice *Hijāma* around 3500 BC, using primitive instruments such as animal horns and bamboo cups. In Arabic civilization, the therapy was known as *Al-Hijāma* (El Sayed, Mahmoud, and Nabo, 2013, p. 111; Ahmadi, Schwebel, and Rezaei, 2008, pp. 37-44).

Historical records also indicate that *Hijāma* was practiced in ancient Macedonia since prehistoric times (3300 BC) for the management of various diseases and health disorders (El Sayed, Mahmoud, and Nabo, 2013; Ahmadi, Schwebel, and Rezaei, 2008, pp. 37-44). Ancient Egyptian civilization recognized *Hijāma* as one of the earliest medical therapies. Evidence of its use dates back to approximately 1550 BC, as documented in the famous Ebers Papyrus. Illustrations depicting cupping instruments have also been identified in the tomb of King Tutankhamun and on the walls of the temple of Kom Ombo, which served as a major medical center during that period (El Sayed, Mahmoud, and Nabo, 2013, p. 111; Ahmadi, Schwebel, and Rezaei, 2008, pp. 37-44; Christopoulou-Aletra, and Papavramidou, 2008, pp. 899-902). In ancient Egypt, *Hijāma* was utilized not only for the treatment of diseases but also for the preservation of general health.

In Traditional Chinese Medicine, cupping therapy has been practiced for more than 2000 years (El Sayed, Mahmoud, and Nabo, 2013, p. 111; Chirali, 1999, pp. 73-86). Its therapeutic application was first systematically documented by Ge Hong (281–341 AD) in the Handbook of Prescriptions for Emergencies (Hanan, and Eman, 2013, pp. 631-642). From Egypt, the practice spread to Greece, where cupping became closely associated with medicine. Renowned Greek physicians such as Hippocrates and Galen strongly advocated the use of *Hijāma*. Hippocrates (460–377 BC) provided detailed descriptions regarding the shape, size, and application of cups, recommending smaller and lighter cups for deep-seated disorders and wider cups for superficial conditions (Mahdavi, 2012, pp. 67-88). Although he mainly promoted dry cupping, he also endorsed limited use of wet cupping for bloodletting purposes.

During the Roman era, physicians such as Celsus and Galen further popularized *Hijāma* and bloodletting therapies for the management of localized and systemic diseases (Turk, and Allen, 2009, pp. 9-15). In Persian medicine, *Hijāma* also occupied an important therapeutic position. Ibn Sina (Avicenna), in his renowned work *Al-Qanoon fi al-Tibb*, described *Hijāma* as a significant therapeutic modality for disease management (Avicenna, 2010, p. 537). The practice of *Hijāma* in Europe and the Middle East evolved alongside humoral theory, which emphasized the balance of the four humours: blood (*dam*), phlegm (*balgham*), yellow bile (*saфра*), and black bile (*sauda*). Prominent physicians, including Galen, Paracelsus, and Ambroise Paré, advocated cupping therapy, which was also commonly practiced by barber surgeons and bath attendants (Ullah, Younis, and Wali, 2007).

In the seventeenth and eighteenth centuries, several European physicians, including Sydenham, Willis, Pitcairn, Mead, and Heberden, continued to recommend *Hijāma* as a therapeutic intervention (Turk, and Allen, 2009, pp. 9-15). In recent decades, there has been growing global interest in *Hijāma* and other complementary and alternative



therapies, particularly in Asia, the Middle East, Europe, and the United States (El Sayed, Mahmoud, and Nabo, 2013, p. 111). In Korean Medicine, cupping therapy is commonly employed to enhance the circulation of qi and alleviate blood stasis through negative pressure and suction techniques. A recent survey among Korean medicine practitioners reported that *Hijāma* is predominantly used for musculoskeletal disorders, particularly involving the neck and shoulder regions (Kim, et al., 2011, p. 146).

Three different treatment modalities are detailed in the traditional textbook of Unāni medicine, namely, '*Ilāj bi'l Tadbīr* (Regimenal therapy), '*Ilāj bi'l Taghdhiya* (Dietotherapy), '*Ilāj bi'l Dawā*' (Pharmacotherapy), and '*Ilāj bi'l Yad* (Surgery). '*Ilāj bi'l Tadbīr* (regimenal therapy) is one of the most popular methods of treatment practiced by ancient Unāni physicians for ages. Regimenal therapy is a modification of *Asbāb Sitta Darūriyya* (six essential factors) and the application of regimens for maintenance of health as well as for disease management. The six essential factors are Air (*Hawā'*), Food and Drink (*Makūl wa Mashrūb*), *Al-Ḥaraka wa'l Sukūn al-Badanī* (bodily movement and repose), *Al-Ḥaraka wa'l Sukūn al-Nafsānī* (psychic movement and repose), *Al-Nawm wa'l Yaqza* (sleep and wakefulness), and *Al-Iḥtibās wa Al-Istifrāgh* (retention and evacuation). Various therapeutic regimens are included under *Ilaj-bit-Tadbeer*, such as *Hijāma* (Cupping), *Ābzān* (Sitz bath), *Hammam* (Bathing), *Dalk* (Massage), *Takmeed* (Fomentation), *Irsale-Alaq* (Leeching), *Fasd* (Venesection), *Ishaal* (Purgation), *Riyazat* (Exercise), *Qai* (Emesis), and *Idrare Baul* (Diuresis). These regimens may be used independently or in combination with other therapeutic modalities. These are essential for every person, and an imbalance in any or all of these factors may lead to various disease conditions. These regimens may be used independently or in combination with other therapeutic modalities (Wani, et al., 2023, pp. 117-132).

The term *Hijāma* is derived from the Arabic word "*Hajm*," meaning "to suck" (Dar, and Lone, 2015). It refers to a therapeutic procedure in which a partial vacuum is created inside cups applied to the surface of the body using heat or mechanical suction. This technique is employed to evacuate morbid materials, divert harmful substances from deeper tissues, and enhance blood circulation at the affected site (Dar, and Lone, 2015; Anjum, et al., 2005, pp. 412-415; Tagil, et al., 2014, pp. 1032-1036). The word *Hijāma* also carries different meanings in Arabic literature, including the removal of scalp hair and the application of cups (Seenghi) (Alam, et al., 2012, pp. 143-148). In the present review, an attempt has been made to elucidate various aspects of *Hijāma* in the Unāni system of medicine.

Materials and Methods

This review study on *Hijāma* was searched in classical Unāni textbooks available in the Kashmir Tibbia College Hospital and Research Centre library, such as *Al-Qanoon Fit Tib*. Vol. 1 & 3 part 2 (Urdu Translation), *Kitabul Hawi*. Vol. 11, Firdous ul Hikmat, *Kaamilus Sana'ah* (Urdu Translation), *Kitab al-Taisir Fil-Mudawat Wal-Tadbeer*, *Kitab Al Mukhtarat Fit Tib*. Vol. 1 & 4, *Matab E Hakim Alwi Khan* (Urdu Translation), *Tibbe Akbar* (Urdu Translation), *Akseere Azam (Al Ikseer)* (Urdu Translation), *Jami-ul-Hikmat*. For a comprehensive review of *Hijāma*, major scientific databases, including PubMed, ScienceDirect, and Google Scholar, were systematically searched. Additionally, the Standard Unani Medical Terminology was consulted to identify appropriate English equiva-



lents for the Unāni terms used throughout the article.

Synonymous

Hijamat, Al-Hijāma, Seenghiyan khichwana, Seenghiyan Lagwana, Pachney lagwana, Sheshay Lagwana and Jiaofa and Baguanfa (Chinese) (Nafisi, 1954, pp. 432-433; Qurshi, 2011, p. 123).

Classification of *Hijāma*

There are various types of cupping as mentioned in Table 1.

Table 1: Types of cupping (Qurshi, 2011, p. 125; Baghdadi, 2007, p. 635; Masihi, 2000, pp. 525-528; Zahrawi, 2012, pp. 702-703; Mohammad, Fasihuzzaman, and Jabeen, 2015, pp. 90-95)

Types	Subtypes
<i>Hijāma bi'l Sharṭ</i> (Wet Cupping)	a. <i>Hijāma Ḍarūriyya</i> (obligatory cupping)
	b. <i>Hijāma Ikhtiyāriyya</i> (Optional)
<i>Hijāma bilā Sharṭ</i> (Dry cupping)	a. <i>Hijāma Mutaharrikah</i> (Gliding or moving cupping)
	b. <i>Hijāma Ghair Mutaharrikah</i> (Static cupping)
	b1. <i>Hijāma bi'l Nār</i> (cupping with fire)
	b2. <i>Hijāma bilā Nār</i> (non-fire cupping)

Conditions of *Hijāma bi'l Sharṭ Ikhtiyāriyya*

Al-Masihi mentioned the following conditions for *Hijāma bish shart ikhtiyari* (Masihi, 2000, pp. 95-96; Ahmedi, and Siddiqui, 2014, pp. 1-14);

1. It should be performed mid-day of the lunar month calendar (Jurjani, 2010, pp. 403-405)
2. It should be performed in sah paher (afternoon) because this time is most suitable for *Hijāma*.
3. It should be performed in the summer. In this season, the Akhlat (humours) become less viscid due to increased Hararat and are easily evacuated through cupping.
4. It should be performed in those persons whose blood is relatively less viscid.
5. It should be performed after the use of syrups having muqawwi-i-mi'da and dafa'-i-mawād properties.
6. It should not be performed in weak and cachexic persons (Qurshi, 2011, p. 127).
7. It should not be performed by anyone under 2 years of age or after 60 years of age (Qurshi, 2011, p. 127; Jurjani, 2010, pp. 403-405; Avicenna, 2010, p. 538).
8. *Hijāma* is not recommended immediately following *Hammām*, because the skin becomes dense and may necessitate deeper incisions to facilitate bloodletting, which can cause considerable pain and physical debility. Moreover, the procedure should be avoided in individuals with thick and viscid blood, especially those who are obese.
9. *Hijāma* should be avoided immediately after sexual intercourse.
10. *Hijāma* should not be performed immediately after strenuous physical activity, as excessive exertion and dehydration may lead to generalized weakness. However, it may



be considered in individuals with a robust physique and thick or viscid blood.

11. *Hijāma* is contraindicated in persons experiencing bodily weakness and laxity resulting from excessive depletion or dissolution of body tissues.

Contraindications for *Hijāma*

1. *Hijāma* should be avoided in individuals suffering from anemia, which may be identified by the physician based on the patient's physical appearance and pulse examination (Nimrouzi, et al., 2014, pp. 128-136).

2. *Hijāma* should preferably be avoided in cold climatic conditions. In unavoidable circumstances, only a small amount of blood may be removed, ideally during the later part of the morning (Nimrouzi, et al., 2014, pp. 128-136).

3. If the causative matter is in abundance, then *Fasd* (venesection) should be done before *Hijāma*. *Fasd* evacuates the morbid material from the body in general, while *Hijāma* bish shart does that from the affected organ in particular (Qurshi, 2011, p. 128).

4. *Hijāmah bi'l Shart* should be avoided in pregnant women, especially during the first five months of gestation, and is also not recommended during menstruation (Mohammad, Fasihuzzaman, and Jabeen, 2015).

5. The procedure should be performed in individuals with a moderate body build, avoiding those who are extremely obese or excessively thin (Qurshi, 2011, p. 128).

6. *Hijāmah bi'l Shart* is recommended only after at least 12 hours have elapsed following intercourse and 3–4 hours after food intake (Nimrouzi, et al., 2014, pp. 128-136).

7. If the aim is to divert the matter from *Shareef uz w* to *adna uz w*, it should be done before the appearance of the peak stage of inflammation (Qurshi, 2011, p. 128).

8. The consumption of certain dietary items, especially eggs, is discouraged for at least 4–6 hours before and after cupping therapy (Nimrouzi, et al., 2014, pp. 128-136).

9. Lean individuals with a hot and dry temperament are advised to take pomegranate juice or chicory with sugar, as this may help prevent the development of unhealthy humours in the stomach. In contrast, other individuals should refrain from eating for approximately three hours after the procedure (Qurshi, 2011, p. 128; Nimrouzi, et al., 2014, pp. 128-136).

Indications of cupping in Unāni Medicine

The therapeutic indications of cupping, categorized according to the type of cupping and anatomical sites, are presented in Tables 2 and 3.

Unani Concept of Mechanism of Action

In the Unāni system of medicine, cupping therapy functions in two ways (Lari, et al., 2017, pp. 21-24; Avicenna, 2006, p. 539; Al-Bedah, et al., 2018, pp. 90-97):

1- *Tanqiya'-i-Mawād* (cleansing of morbid matter)

In Unani literature, disease is considered to arise from an imbalance of the *Akhlāṭ* (humours) leading to the accumulation of morbid material at specific sites within the body. *Hijāma bi'l Shart* acts on the principle of *Tanqiya'-i-Mawād* and hereby facilitates the removal of morbid substances from the affected region (Lari, et al., 2017, pp. 21-24; Avicenna, 2006, p. 539). Jālinus stated that when the humours become dense and



Table 2: Indications of *Hijāma bi'l Sharṭ* (wet cupping) according to anatomical sites (Masihi, 2000, pp. 95-96; Jurjani, 2010, pp. 403-405; Avicenna, 2010, p. 540; Kim, et al., 2011, p. 146; Arzani, 2002, pp. 138-140)

S. Number	Anatomical Sites	Indications
1	<i>Al-Hijāma 'ala'l Hāma</i> (sinciput)	Delirium, vertigo, pruritus of the eye, iridoptosis
2	<i>Al-Hijāma 'ala'l Qamaḥduwa</i> (Occiput)	Delirium, vertigo, pruritus of the eye, iridoptosis, Mania, Giddiness (Masihi, 2000, pp. 95-96; Jeelani, 1996, pp. 185-187)
3	<i>Al-Hijāma 'ala'l Nuqra</i> (Nape of the neck)	Headache, conjunctivitis, meningitis and stomatitis, otalgia, melasma, freckles, naevus, blepharitis, pruritus of the eye, halitosis (Anees, et al., 2015, pp. 88-100)
4	<i>Al-Hijāma 'ala'l Akhda'ayn</i> (cupping over sides of neck)	Tremors, glossalgia, gingivitis, conjunctivitis, otitis, headache, migraine, diphtheria
5	<i>Al-Hijāma Taḥt al-Dhaqan</i> (cupping below chin)	Mouth ulcers, stomatitis, tonsillitis, glossitis, oral thrush, and halitosis (Masihi, 2000; Arzani, 2002, pp. 138-140)
6	<i>Hijāma al-Udhunayn</i> (cupping on ears)	Blepharitis, headache
7	<i>Al-Hijāma 'ala'l Kāhil</i> (Inter scapular region)	Bronchial asthma, congestion, sanguineous diphtheria, frozen shoulder/shoulder pain, hemoptysis, sanguineous diseases of the thorax
8	<i>Hijāma al Mankib</i> (Shoulders)	Palpitation due to heat and congestion, liver diseases (right mankib), spleen diseases (left mankib), tertian fever
9	<i>Hijāma Baṭṭayn al-Zandayn</i> (Ventral aspect of forearm and wrist)	Sanguineous chronic pruritis, fissure in hand, carpal tunnel syndrome
10	<i>Nāghis</i> (lateral to the spine, near the liver and spleen)	Hepatalgia (right naghiz), splenalgia (left naghiz)
11	<i>Al-Hijāma 'ala'l Thadyayn</i> (Below mammary glands/ bust line)	Menorrhagia, metrorrhagia, puerperal diseases, and epistaxis (Lari, et al., 2017, pp. 21-24)
12	<i>Warkayn</i> (Buttocks)	Sciatica and dislocation of the hip, hemorrhoids, metritis, proctitis, haematuria, burning micturition, anal throbbing, oophoritis, and boils of the thigh (Lari, et al., 2017, pp. 21-24)
13	<i>Al-Hijāma 'ala'l Qaṭan</i> (Lumbar region)	Boils, pustules, and pruritus of the thigh, gout, hemorrhoids, filariasis, and pruritus of the thigh
14	<i>Hijāma al-'Uṣ'us</i> (Coccyx)	Piles, per rectal bleeding, oophoritis, pruritus vulvae
15	<i>Hijāma al-Rukba</i> (Knee)	Swelling of the knee, sanguineous arthritis, infected wounds of the thigh (Anees, et al., 2015, pp. 88-100)
16	<i>Al-Hijāma 'ala'l Maq'ada</i> (cupping near anal orifice)	Intestinal disorders, piles, a disorder of the anus, diversion of matter from the head, and menstrual disturbances
17	<i>Al-Hijāma 'ala'l Fakhid-hayn</i> (Thighs)	Orchitis, leg ulcers and metritis, uterine hemorrhage, hemorrhoids, fissure in ano (Zaidi, et al., 2016, pp. 81-94)
18	<i>Al-Hijāma 'ala'l Sāq</i> (Shank/Calf muscles)	Chronic knee arthritis, chronic cystalgia, Renal colic, chronic metritis, and amenorrhea (Zaidi, et al., 2016, pp. 81-94)
19	<i>Mahajim Arqubeen</i> (Tendo-calcaneus or tendo achilis)	Chronic knee arthritis, chronic cystalgia, Renal colic, chronic metritis, amenorrhea
20	<i>Al-Hijāma 'ala'l Ka'bayn</i> (Ankle joints)	Amenorrhea, sciatica, gout

stagnant within the joints, *Hijāma* proves particularly beneficial in dispersing and evacuating these accumulated materials. It increases blood circulation, dilates the cutaneous pores, improves tissue nourishment with the supply of fresh blood, and strengthens the body's eliminative. Through these actions, it assists in the evacuation of *Akhlāt-i-Fāsida*



(morbid humours) and helps restore humoral equilibrium by eliminating deranged qualities such as *Hār* (heat), *Bārid* (cold), *Raṭb* (moistness), and *Yābis* (dryness) (Lari, et al., 2017, p. 21-24).

2- *Imāla--i-Mawād* (diverting morbid matter)

Hijāma bilā Sharṭ (Dry Cupping) operates on the principle of *Imāla-i-Mawād* (Diversion). This technique involves moving morbid material from one location to another. Through this process diversion, *Hijāma bilā Sharṭ* enables *Ṭabī'at Mudabbira-i-Badan* to resolve the morbid matter. *Hijāma bilā Sharṭ* helps relieve congestion, resolve obstructions, and restore the normal flow of blood circulation, thereby reducing muscular tension and pain. It is also believed to divert inflammation and pressure from the deeper vital organs, particularly the heart, brain, lungs, liver, and kidneys, towards the skin and superficial tissues of the body.

As a result, the immune system is strengthened, and the healing process is facilitated, promoting the body's optimal functioning (Lari, et al., 2017, pp. 21-24; Avicenna, 2006, p. 540).

Table 3: Indications of *Hijāma-bila-shart* (Dry Cupping) according to anatomical sites (Masihi, 2000, pp. 95-96; Jurjani, 2010, pp. 403-405; Avicenna, 2010, p. 288; Nimrouzi, et al., 2014, pp. 128-136; Zahrawi, 2012, pp. 322-324; Avicenna, 2006, p. 539; Nayab, Anwar, and Quamri, 2011, pp. 697-701; Mehta, and Dhapte, 2015, pp. 127-134)

S. Number	Sites	Indications
1	Below the mammary glands	Metrorrhagia, menorrhagia, and epistaxis
2	Around the umbilical region	Colicky abdominal pain, dyspepsia, painful menstruation, tenesmus
3	Over the abdomen, especially the epigastrium	Epistaxis, diarrhea, acne
4	Lower abdomen	Inguinal hernia
5	Hypochondriac region	Hepatic problems (right side), Spleen problems (left side) due to riyah
6	kidney	Kidney stones
7	Over hip	sciatica
8	Anus	Intestinal and menstrual disorders, hemorrhoids
9	Posterior area of thigh	Sciatica
10	On the 1st cervical vertebrae	Facial paralysis
11	Calf muscles	Diphtheria (Qamri, 2008, pp. 237-239)
12	On chest ribs and calf muscles	Pneumonia (Qamri, 2008, pp. 237-239)
13	On the chest, between the shoulders	Pleurisy (Qamri, 2008, pp. 237-239)
14	Stomach, the site of liver	Syncope (Qamri, 2008, pp. 237-239)

Mechanism of action in modern medicine

Although the exact therapeutic basis of *Hijāma* has not yet been fully elucidated, several theories have been suggested to explain its mode of action and the diverse effects produced by cupping therapy. The cupping session is linked to biological and mechanical



processes, according to several researchers. (Lari, et. al., 2017, pp. 21-24; Al-Bedah et. al., 2018, pp. 90-97; Anees et.al., 2015, pp. 88-100) (Table 4).

Pain Gate Theory: According to this concept, pain impulses are carried from the affected area to the brain through specific pathways. The application of a cupping cup creates another sensory stimulus that interferes with these pain impulses, thereby reducing the perception of pain through the same pathway. (Lari, et al., 2017, pp. 21-24; Al-Bedah, et al., 2018, pp. 90-97; Anees, et al., 2015, pp. 88-100).

Reflex Zone Theory in Cupping Therapy: According to this theory, dysfunction in an internal organ produces reflex changes on the corresponding area of the skin, which may manifest as redness, tenderness, or swelling. When cupping is applied, the cutaneous receptors are stimulated, leading to improved blood circulation and enhanced blood supply to both the skin and related internal organs through neurovascular reflex mechanisms (Lari, et al., 2017, pp. 21-24; Al-Bedah, et al., 2018, pp. 90-97; Anees, et al., 2015, pp. 88-100).

Prostaglandin theory: Inflammation in the body leads to the production of prostaglandins, which play an important role in transmitting pain impulses to the brain. *Hijāma bi'l Sharṭ* is believed to help eliminate these inflammatory mediators from the body, thereby contributing to pain relief (Lari, et al., 2017, pp. 21-24; Al-Bedah, et al., 2018, pp. 90-97; Anees, et al., 2015, pp. 88-100).

Endorphins and enkephalin production theory: These are endogenous pleasure compounds, which are natural substances produced by our bodies and which reduce pain. 91,99

Nitric Oxide Theory: According to this theory, nitric oxide is an important biological mediator released in response to tissue injury and is also believed to be produced during or after *Hijāma bi'l Sharṭ* (wet cupping). Nitric oxide exerts several beneficial effects, including vasodilatation, improvement of blood circulation, relaxation of smooth muscles leading to relief of spasms, anti-thrombotic activity, and anti-inflammatory action. It also helps prevent narrowing of blood vessels, thereby promoting better vascular function (Lari, et al., 2017, pp. 21-24; Al-Bedah, et al., 2018, pp. 90-97; Anees, et al., 2015, pp. 88-100).

Taibah theory: According to Taibah theory, the skin is raised when negative pressure (suction force) is exerted using cups on the skin's surface because the skin is elastic. The gradual expansion during cupping reduces local capillary pressure according to Boyle's law, leading to increased capillary filtration and accumulation of lymphatic and interstitial fluids in the skin. This decreases pain by dilating chemical substances, inflammatory mediators, and nociceptive substances, bathing nerve endings in gathered fluids, and rupturing tissue adhesions (Lari, et al., 2017, pp. 21-24; Al-Bedah, et al., 2018, pp. 90-97; Anees, et al., 2015, pp. 88-100). According to the *Taibah* theory, *Hijāma bi'l Sharṭ* is considered a minor excretory surgical procedure that facilitates the removal of pathological substances from the body, in a manner comparable to renal glomerular filtration and abscess drainage. (Lari, et al., 2017, pp. 21-24; Al-Bedah, et al., 2018, pp. 90-97; Anees, et al., 2015, pp. 88-100).

Chinese Theory: According to practitioners of Traditional Chinese Medicine, diseases arise due to stagnation or blockage in the flow of *Qi* (vital energy). Cupping therapy is believed to restore the normal flow of *Qi* by removing these obstructions and re-establishing the body's natural balance (Lari, et al., 2017, pp. 21-24; Al-Bedah, et al., 2018,



pp. 90-97; Anees, et al., 2015, pp. 88-100).

Analgesic effects of cupping therapy: The mechanical stimulation and minor skin injury produced during wet cupping therapy may activate $A\beta$ nerve fibers and inhibitory neural pathways at the spinal cord level, thereby modulating pain perception. The superficial skin incisions and negative pressure created during the procedure draw out a small quantity of blood and generate a nociceptive stimulus, which may activate diffuse noxious inhibitory control mechanisms. This response is believed to reduce chronic musculoskeletal pain, including its emotional and affective components, through modulation of limbic system activity. Therefore, the analgesic effect of wet cupping therapy may be attributed to both tactile and nociceptive stimulation produced during the procedure (Lari, et al., 2017, pp. 21-24; Al-Bedah, et al., 2018, pp. 90-97; Anees, et al., 2015, pp. 88-100).

Increasing blood circulation theory: Cupping therapy is believed to improve blood circulation, relieve congestion, and facilitate the removal of inflammatory substances and toxins from the tissues, thereby helping to reduce inflammation and pain (Lari, et al., 2017, pp. 21-24; Al-Bedah, et al., 2018, pp. 90-97; Anees, et al., 2015, pp. 88-100).

Table 4: Mechanism of Action and Therapeutic Effects of Cupping Therapy

Theories	Effects on the body
<i>Pain gate theory</i>	Pain reduction effect
<i>Diffuse noxious inhibitory mechanism</i>	Pain reduction effect
<i>Theory of Reflex Zone Stimulation</i>	Pain reduction effect
<i>Nitric Oxide Release Theory</i>	Anti-inflammatory and improved blood circulation
Theory of Immune Modulation	Immune regulatory effect
Blood circulation theory	Hematological effect

Scientific studies on cupping therapy

Several clinical studies and case-based investigations have highlighted the therapeutic potential of cupping therapy across diverse clinical conditions. Kouser, et al., (2024) demonstrated that wet cupping significantly improves symptoms of cervical spondylosis. Similarly, Begum et al., (2023) reported that dry cupping, when used adjunctively with Unani medicine, is effective in managing primary dysmenorrhea. In another study, Khatoon et al., (2024) found that wet cupping offers notable clinical benefits in patients with rheumatoid arthritis.

Research by Kalam, Suhail, and Ibrahim, (2024) further supports the efficacy of cupping modalities, showing that wet cupping combined with Unani therapeutic measures is beneficial in chronic migraine. In a separate investigation, Kalam, Jalal, and H, (2024) also reported the effectiveness of dry cupping in improving functional outcomes in patients with frozen shoulder.

Conclusion

Hijāma is a significant therapeutic modality of *'Ilāj bi'l Tadbīr* in the Unani system of medicine that primarily functions through the evacuation and diversion of morbid substances from the body. Contemporary studies indicate that *Hijāma* may contribute to



health promotion and disease management, serving both preventive and therapeutic purposes. It has demonstrated potential benefits in a variety of disorders, particularly those related to the musculoskeletal, reproductive, circulatory, and nervous systems.

Despite its growing acceptance in complementary and integrative medicine, further scientific validation is required to establish its efficacy according to modern medical standards. The development of standardized operating procedures (SOPs), evidence-based guidelines, and methodological research protocols is therefore essential. Future investigations involving large sample sizes, randomized controlled trials, and long-term follow-up conducted by qualified clinicians and researchers are strongly recommended to evaluate the clinical effectiveness and safety of *Hijāma* in diverse disease conditions.

Acknowledgements

We are thankful to the library staff of Kashmir Tibbia College Hospital and Research Centre for their support in compiling this work.

Authors' Contribution

Khalid Rahim Wani conceptualized and curated the study plan, conducted the literature review, analyzed the collected information, and drafted the original manuscript. Abdul Nasir Ansari assisted in the literature survey, contributed to the manuscript revision, and provided critical editorial input. Mohd Nayab performed the final review and proofreading of the manuscript. All authors read and approved the final version of the manuscript.

Funding

None.

Conflict of Interest

None.

References

- Ahmadi, A., Schwebel, DC., and Rezaei, M., 2008. The efficacy of wet-cupping in the treatment of tension and migraine headache. *Am J Chin Med*, 36, pp. 37-44.
- Ahmedi, M., and Siddiqui, MR., 2014. The value of wet cupping as a therapy in modern medicine-An Islamic Perspective. *Webmedcentral*, 5(12), pp. 1-14. Doi:10.9754/journal.wmc.2014.004785.
- Alam, M., Mannan, A., Siddique, M., and Ilahi, A., 2012. Hijamah (cupping): A Greco-Islamic Perspective. *J Int S hist Islamic Med*, 10(11), pp. 143-8.
- Al-Bedah, A.M.N., Elsubai, I.S., Qureshi, N.A., Aboushanab, T.S., Ali, G.I.M., and El-Olemy, A.T., 2018. The medical perspective of cupping therapy: Effects and mechanisms of action. *J Traditional and Complementary Medicine*, 9(2), pp. 90-7.
- Anees, S., Arafath, Y., Naaz, A., and Khan, M.Q., 2015. Hijamah (Cupping Therapy) as a preventive medicine- a retrospective analytical study. *Int J AYUSH*, 4(2), pp. 88-100.
- Anjum, N., Jamil, S., Hannan, A., Akhtar, J., Ahmad, B., 2005. Clinical efficacy of Hijamat (Cupping) in Waja-ul-Mafasil (Arthritis). *Indian J Tradit Knowl*, 4, pp. 412-415.
- Arzani, A., 2002. *Meezan -ut-Tibb*. New Delhi: Idara Kitab us Shifa.
- Avicenna, I., 2006. *Kulliyat-i-Qanoon*. Translated from Urdu by Hkm Kabiruddin. New Delhi: Ejaz Publishing House.



- Avicenna, I., 2010. *Al-Qanoon Fit Tib*. Vols. I & III. Translated from Urdu by G.H. Kantoori. New Delhi: Idara Kitab us Shifa.
- Baghdadi, I.H., 2007. *Kitab al-Mukhtarat fil-Tib*. Part IV. 1st ed. Central Council For Research In Unani Medicine. New Delhi: Ministry of Health and Family Welfare, Govt. of India.
- Begum, A., Sultana, S., and Sultana, A., 2023. An open-labelled randomized comparative study of Unani formulations and Hijama Bila Shart in Tashanuji Ushr al-Tamth. *Pharmacognosy Research*, 16(1), pp. 183–190. doi: 10.5530/pres.16.1.23.
- Chirali, I.Z., 1999. *The cupping procedure*. Churchill Livingstone, London: Traditional Chinese Medicine Cupping Therapy.
- Christopoulou-Aletra, H., and Papavramidou, N., 2008. Cupping: an alternative surgical procedure used by Hippocratic physicians. *J Altern Complement Med*, 14, pp. 899-902.
- Dar, A.K., and Lone, A.H., 2015. Therapeutic application of Al-Hijamah (cupping therapy) in osteoarthritis of the knee. *Int J Research and Development in Pharmacy & Life Science*, 4(3), pp. 1540-1544.
- El Sayed, S.M., Mahmoud, H.S., and Nabo, M.M.H., 2013. Methods of Wet Cupping Therapy (Al-Hijamah): In Light of Modern Medicine and Prophetic Medicine. *Altern Integ Med*, 2, p. 111. doi:10.4172/2327-5162.1000111
- Hanan, S.A., and Eman, S.E., 2013. Cupping Therapy (Al-Hijāma): Its Impact on Persistent Non-Specific Lower Back Pain and Client Disability. *Life Sci J*, 10(4s), pp. 631-642.
- Jeelani, G., 1996. *Makhzanul Hikmat*. Vol. II. New Delhi: Aijaz Publishing House.
- Jurjani, A.H., 2010. *Zakheera Khwarzam Shahi*. New Delhi: Idara Kitab-US-Shifa.
- Kalam, A., Jalal, B., and H, S., 2024. Effect of Hijama Bila Shart (Dry cupping) and Unani formulation on Waja-i-Mafsal-i-Katif (Frozen Shoulder): A case study. *Int J AYUSH Case Reports*, 8(3), pp. 351–356.
- Kalam, A., Suhail, S.B., and Ibrahim, T., 2024. Effect of Hijama bi'l Shart (Wet cupping) with Unani formulation in a patient of Shaqīqa-i-Muzmin (Chronic Migraine): A case report. *Int J AYUSH Case Reports*, 8(4), pp. 537–544. doi: 10.70805/ija-care.v8i4.654.
- Khatoun, S., Shoaib, M., Azmat, J., and Anwar, M., 2024. Evaluation of the efficacy of Unani formulation along with Hijama bi'l Shart in patients of rheumatoid arthritis. *Int J Unani and Integrative Medicine*, 8(2), pp. 111–118. doi: 10.33545/2616454X.2024.v8.i2b.287.
- Kim, J-I., Kim, T-H., Lee, MS., Kang, JW., Kim, KH., Choi, J-Y., et al., 2011. Evaluation of wet-cupping therapy for persistent non-specific low back pain: A randomized, waiting-list controlled, open-label. *Parallel-Group Pilot trial*, 12(1), p. 146. doi: 10.1186/1745-6215-12-146
- Kouser, V.H., Nayab, M., Ansari, A.N., Mir, M.A., and Anwar, M., 2024. Efficacy of wet cupping in the pain management of cervical spondylosis - A randomized, controlled, open-label, parallel-group clinical trial. *J bodywork and movement therapies*, 39, pp. 126–131. https://doi.org/10.1016/j.jbmt.2024.02.044
- Lari, A., Nayab, M., Tausif, M., Lari, J., Alam, A., 2017. Therapeutic potentials of hijama bila-shart (dry cupping therapy): a review. *Int J Unani Integ*, 1(1), pp. 21–24.
- Mahdavi, M.R.V., 2012. Evaluation of the Effects of Traditional Cupping on the Biochemical, Hematological, and Immunological Factors of Human Venous Blood. A Compendium of Essays on Alternative Therapy, Rijeka, Croatia: InTech.
- Masihi, I.Q., 2000. *Kitab Al-Umda Fil Jarahat*. Vols. 1 & 2. New Delhi: CCRUM, Ministry of Health and Family Welfare, Govt. of India.
- Mehta, P., and Dhapte, V., 2015. Cupping therapy: A prudent remedy for a plethora of medical



- ailments. *J Traditional and Complementary Medicine*, 5(3), pp. 127–34.
- Mohammad, Sh.H., Fasihuzzaman, and Jabeen, A., 2015. Hijamah (Cupping therapy): A noble method of treatment in Unani medicine. *Int J Res Ayurveda Pharm*, 6(2), pp. 207-215.
- Nafisi, B., 1954. *Tarjuma Wa Sharah Kulliyate Nafisi*. Translated from Urdu by M. Kabeeruddin. New Delhi: Idara Kitab us Shifa.
- Nayab, M., Anwar, M., and Quamri, M.A., 2011. Clinical study on Waja-ul-Mafasil and evaluation of the efficacy of hijamat-bila-shurt in the treatment. *Indian J Tradit Knowl*, 10(4), pp. 697–701.
- Nimrouzi, M., Mahbodi, A., Jaladat, A.M., Sadeghfard, A., Zarshenas, M.M., 2014. hijamat in traditional Persian medicine. *J Evidence-Based Complementary & Alternative Medicine*, 19(2), pp. 128–36.
- Qamri, A.M., 2008. *Minhaj Ul Ilaj*. Translated from Urdu by Ghina Muna. New Delhi: CCRUM.
- Qurshi, H.M., 2011. *Jami-ul-Hikmat*. New Delhi: Idara Kitab us Shifa.
- Tagil, S.M., Celik, H.T., Ciftci, S., Kazanci, F.H., Arslan, M., Erdamar, N., et al., 2014. Wet-cupping removes oxidants and decreases oxidative stress. *Complementary Therapies in Medicine*, 22(6), pp. 1032–6.
- Turk, J.L., and Allen, E., 2009. History of medicine: Bleeding and cupping. *Annals of the Royal College of Surge herapies of medicine*, pp. 9-15.
- Ullah, K., Younis, A., and Wali, M., 2007. An investigation into the effect of Cupping Therapy as a treatment for Anterior Knee Pain and its potential role in Health Promotion. *The Internet J Alternative Medicine*, 4(1).
- Wani, K.R., Khan, F., Nayab, M., and Ansari, AN., 2023. Understanding The Concept, Classifications, and Effects of Dalk (Massage Therapy) In Unani Medicine and Contemporary Era. A Review. *J Emerging Technologies and Innovative Research*, 10(4), pp. 117-132.
- Zahrawi, A.Q., 2012. *Jarahiyat zahrawi*. Translated from Urdu by Hk. Nasir Ahmad Kakorvi. New Delhi: CCRUM, Ministry of Health and Family Welfare, Govt. of India.
- Zaidi, S.M., Jameel, S.S., Jafri, K., Khan, S.A., and Ahmad, E., 2016. Ilaj bil Hijamah (cupping therapy) in Unani System of Medicine: Anecdotal practice to evidence-based therapy. *AMHA-Acta medico-historica Adriatica*, 14(1), pp. 81-94.

